



Two weeks prior to your surgical date, **DO NOT** take any Aspirin or any medication containing Aspirin. Aspirin (salicylate) can cause an abnormality of blood clotting. This could cause excessive bleeding at the time of surgery. Tylenol (acetaminophen) may be used for pain.

LISTED BELOW ARE SOME OF THE DRUGS WHICH CONTAIN ASPIRIN

Alka Seltzer	Buff-A comp #3 tablets	Empirin with codeine	Oxycodone terephthalate & aspirin tabs
A.P.C. with codeine tabloid brand	Bufferin	Equagesic	Pepto-Bismol
Anacin analgesic capsules or tablets	Bufferin with codeine #3	Excedrin Extra strength	Percodan & Percocan – demi tablets
Anacin maximum strength analgesic tablets	Cama Arthritis Pain-Reliever	Excedrin Migraine	Ponstel (mefenamic acid)
Arthritis Bayer timed release aspirin	Celebrex	Extra strength Bufferin	Propoxyphene compound 65
Arthritis pain formula made by the makers of Anacin analgesic tablets	Congespirin	Capsules & tabs	Relafen (nabumetone)
Arthritis Strength Bufferin	Cosprin	4-Way cold tablets	Robaxial tablets
Ascriptin	Cosprin 650	Fiorinal	Sk-Oxycodone with aspirin tablets
Ascriptin A/D	Darvon with A.S.A.	Fiorinal with codeine	Sk-65 compound capsules
Ascriptin with codeine	Darvon-N with A.S.A.	Germisyn tabs and power	Soma Compound (carisoprodol)
Aspergum	Dayori (oxaprozin)	Halfprin	Stanback Powder
Aspirin suppositories	Dewitts	Hycy-Pap	Supac
Aspirin with codeine tablets	Digesic	Lodine (etodolac)	Synalgos – DC capsules
Aspirin 325 mg with codeine tablets	Dihydrocodeine compound tablets	Maximum Bayer Aspirin	Talwin Compound
Axotal	Disalcid (salsalate)	Mepro-compound tablets	Vanquish
Azulfidine (sulfasalazine)	Doans	Methocarbamol with aspirin tablets	Vemin
Bayer aspirin	Dolobid (diflunisal)	Midol	Vicoprofen (ibuprofen + hydrocodone)
Bayer children’s chewable aspirin	Double-A tablets	Mobic (meloxicam)	Vioxx
Bayer children’s cold tablets	Easprin	Mobigesic	Voltaren (diclofenac sodium)
BC Powder	Ecotrin Enteric coated aspirin	Momentum	Zorpri
Buff-A comp tablets	Ecotrin maximum strength tablets	Norgesic and Norgesic Forte	
	Ecotrin regular strength safety coated aspirin tablets	Oxycodone Hydrochloride	

TOPICAL SALICYLATES:

Arthritis	Bengay	Myoflex	Sportscreme
Aspercreme	Mobisyl	Painbuster	Theragesic

THE FOLLOWING MEDICATIONS ALSO PROLONG BLEEDING TIME AND SHOULD BE AVOIDED

TWO WEEKS PRIOR TO SURGERY:

Acthar	Butazolidin	Meclomen	Naproxen	Oxalid
Advil & Aleve	Clinoril	Motrin	Nuprin	Plaquenil
Anaprox	Feldene	Nalfon	Orabase	Rufen
Axolid	Indocin	Naprosyn	Ascalgesic	Tanderil

BLOOD THINNERS:

Coumadin (warfarin)	Ticlid (ticlopidine)	Plavix (clopidogrel)	Plavix (clopidogrel)
---------------------	----------------------	----------------------	----------------------

NSAIDS:

- Ansaïd (flurbiprofen)
- Arthrotec (diclofenac + misoprostol)
- Bextra
- Cataflam (diclofenac potassium)

ENT Medical Center
5258 Dijon Drive
Baton Rouge, Louisiana 70808
(225) 769-1090
1-800-960-1 ENT